


DNA metabarcoding for environmental biodiversity evaluation

Catalina Stoica, Daniela Ionica, Lucian Ionescu, Daniel Mitru, Alina Roxana Banciu, Stefania Gheorghe, Irina Lucaciu, Mihai Nita-Lazar

National Research and Development Institute for Industrial Ecology – ECOIND, 71-73 Drumul Podu Dambovitei, district 6, 060652, catalina.stoica@incdecoind.ro, Romania


Introduction

The decline of biodiversity and aquatic ecosystem functioning is triggered by human activities which generate pollution, habitat degradation, flow modification, overexploitation and the spread of invasive species as well as climate change (Fig.1). Various legislation documents were globally adopted, including Water Framework Directive (WFD, 2000/60/EC) at European level, to reduce the degradation of aquatic ecosystems. The WFD have confirmed that the focal point of ecosystem integrity should be based on its biology rather than its chemical and physical characteristics.



Aquatic ecosystem degradation

At the beginning of the 21th century, numerous assessment methods were proposed to focus on the biology as proposed in the WFD objectives outlined by the Member States of European Union. Most of the methods relied on morphological taxonomy and comprised similar workflow steps for aquatic bioassessment, such as: i) sampling of “biological quality elements” (BQE); ii) sorting; iii) identification morphology-based; iv) drawing-up the taxa list; v) assessment to an ecological quality class.



The DNA metabarcoding approach is a viable alternative to morphological taxonomy methods and it is gradually integrated in the environmental biodiversity assessment studies. eDNA metabarcoding workflow includes: i) isolation of eDNA from a single environmental component (water, sediment, soil) or bulk samples; ii) PCR amplification of a marker gene targeting biotic communities; iii) high-throughput sequencing of obtained amplicons; iv) quality filtering of identical sequences to obtain Individual Sequence Units (ISU); v) clustering of ISU into Operational Taxonomic Units (OTUs); vi) taxonomic assignment of OTUs and further data analysis-index calculation and quality assessment.

Acknowledgments

This work was financially supported by NUCLEU Program, project no. PN 19-04 02 01.

The advantages of using eDNA metabarcoding over morphological taxonomy methods requires

- cost-effective and reduced time of analysis;
- high accuracy of species detection;
- the use of bulk species without a priori knowledge;
- early life stages, partially destroyed or fragmented species;
- increase comparability across geographic regions.

In addition, eDNA metabarcoding could be integrated in monitoring studies across various ecosystems, such as freshwater, terrestrial, marine, estuarine.

eDNA metabarcoding applicability

Ecosystem type			
Freshwater	Terrestrial	Marine	Estuarine
Water / Sediment	Soil	Water / Sediment	Water / Sediment
Microbes	Fungi	Microbes	Microbes
Diatoms	Microbes	Microbes	Microbes
Plankton	Parasitoids	Plankton	Plankton
Plankton	Arthropods	Fungi	Fungi
Protists	Plants	Protists	Protists
Plants	Vertebrates	Diatoms	Diatoms
Fungi		Plants	Plants
Invertebrates		Invertebrates	Invertebrates
Vertebrates		Vertebrates	Vertebrates

Conclusions

This study focused to pinpoint the usefulness of DNA metabarcoding for environmental biodiversity assessment compared with the traditional morphology based method. Although eDNA method has been developed and largely used in the last decade, there are several limitations that need to be consider in the future researches, such as:

- drawing-up standardized molecular protocols;
- harmonization of DNA preservation and isolation methods;
- selection of DNA indexes and PCR primers;
- the reference database for bioindicator taxa is incomplete;
- development of new biotic indices based on metabarcoding results as well as taxonomic assignment of OTUs and clustering.

References

- [1] Dudgeon, D., Arthington, A.H., Gessner, M.O, Kawabata, Z.I., Knowler, D.J., Leveque, C., Naiman, R.J., Prieur-Richard, A.H., Soto, D., Stiassny, M.L.J, Sullivan, C.A., 2006, *Biol. Rev.* 81, 163-182.
- [2] Leese, F., Bouchez, A., Abarenkov, K., Altermatt, F., Borja, A., Bruce, K., Ekrem, T., Ciampor, Jr., F., Ciamporova-Zatovicova, Z., Costa, F.O., Duarte, S., Elbrecht, V., Fontaneto, D., Franc, A., Geiger, M.F., Hering, D., Kahlert, M., Strojil, B.K., Kelly, M., Keskin, E., Liska, I., Mergen, P., Meissner, K., Pawlowski, J., Penev, L., reyjol, Y., Rotter, A., Steinke, D., van der Wal, B., Vitecek, S., Zimmermann, J., Weigand, A.M., 2018, *Advan. Ecol. Research*, 58, 63-99.
- [3] Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., Willerslev, E., 2012, *Mol. Ecol.* 21, 2045-2050.
- [4] Ruppert, K.M., Kline, R.J., Rahman, M.S., 2019, *Glob. Ecol. Conserv.*, 17, e00547.
- [5] Pawlowski, J., Kelly-Quinn, M., Altermatt, F., Apotheloz-Perret-Gentil, L., Beja, P., Boggero, A., Borja, A., Bouchez, A., Cordier, T., Domaizon, I., Joao Feio, M., Filipe, A.F., Fornaroli, R., Graf, W., Herder, J., van der Hoorn, B., Jones, J.I., Sagova-Mareckova, M., Moritz, C., Barquin, J., Piggott, J.J., Pinna, M., Rimet, F., Rinkevich, B., Sousa-Santos, C., Specchia, V., Trobajo, R., Vasselon, V., Vitecek, S., Zimmerman, J., Weigand, A., Leese, F., Kahlert, M., 2018, *Sci. Total Environ.*, 637-638, 1295-1310.